


## CLENPIQ SPLIT DOSING INSTRUCTIONS

---

### CLENPIQ REGIMEN:

Your kit should come with the following items:

(2) 5-ounce bottles of Laxative Liquid

<p>PART ONE – THE EVENING BEFORE:</p> <ol style="list-style-type: none"><li>1) Drink one bottle of CLENPIQ directly from the bottle. Do not mix or dilute the solution.</li><li>2) Drink five (5) 8-ounce cups of clear liquid, taken at your own pace within the next five hours.</li></ol>	
<p>PART TWO – THE MORNING OF PROCEDURE</p> <ol style="list-style-type: none"><li>1) Drink one bottle of CLENPIQ directly from the bottle. Do not mix or dilute the solution.</li><li>2) Drink four (4) 8-ounce cups of clear liquid. Finish these four hours prior to your procedure time.</li></ol>	

### **IMPORTANT REMINDERS!**

- PATIENTS TAKING GLP 1 AGONISTS ARE REQUIRED TO COMPLETE THEIR PREP A FULL FOUR HOURS PRIOR TO THE PROCEDURE, HAVING NOTHING BY MOUTH WITHIN THIS FOUR-HOUR WINDOW.
- IF YOU FEEL YOU ARE UNABLE TO COMPLETE YOUR PREP WITHIN THE ALLOTTED TIME, YOU MAY START YOUR MORNING DOSE ONE HOUR EARLIER.
- DO NOT TAKE THE DOSE THE NIGHT BEFORE AND THE DOSE THE MORNING OF THE PROCEDURE WITHIN EIGHT HOURS OF EACH OTHER.
- YOU WERE GIVEN A PRESCRIPTION FOR ONDANSETRON; THIS MEDICATION CAN BE TAKEN TO HELP WITH NAUSEA DURING YOUR PREP. TAKE 1-2 PILLS APPROXIMATELY 10-15 MINUTES PRIOR TO PREP DOSE.